

## **NSAIDs Till Prompt**

## Safety messages for non-steroidal anti-inflammatory drugs

Drug name	Some common brand names	
Aspirin	Anadin, Disprin	
Celecoxib		
Diclofenac		
Etodolac		
Flurbiprofen		
Ibuprofen	Nurofen,	
	Some cold/flu treatments	
Indometacin		
Mefenamic Acid		
Meloxicam		
Nabumetone		
Naproxen	Feminax Ultra	
Piroxicam		
Sulindac		

Keep beside the till - please see overleaf for safety messages.







## **NSAIDs Safety Messages**

Message	Why?	Any other information?
Always take this medicine with or after food.	Helps to reduce or avoid gastrointestinal side effects.	
Tell us if you get any side effects.	Earlier recognition of side effects allows action to be taken before they develop into something more serious.	<ul> <li>Side effects to look out for:</li> <li>nausea, vomiting, stomach pain or acid, black stools.</li> <li>What if the patient reports side effects?</li> <li>If a patient reports vomiting blood or black stools, this should be referred to a GP urgently as it indicates bleeding in the gastrointestinal tract.</li> <li>If a patient reports nausea, stomach pain or acid, it would be appropriate to stop the NSAID temporarily to see if this improves the symptoms.</li> </ul>
Be aware of the medicine sick day rules.	Some medicines, including NSAIDs, should be stopped temporarily during dehydrating illness.  This is because continuing to take them when dehydrated increases the risk of serious adverse events, in particular acute kidney injury.	Explain that dehydration can occur with vomiting and diarrhoea, therefore the NSAID should be stopped until the patient is fully recovered. It can then be restarted. Explain that these rules are to cover situations like sickness bugs and food poisoning: it is separate from the advice above on gastrointestinal side effects.  For patients who are buying a single pack of NSAIDs for a single episode of pain like a headache, there is no need to explain the medicine sick day rules. The rules are for people who take NSAIDs every day.